

**List of the delphacid planthoppers of Central and South America, including the Caribbean.**

**Draft Feb. 3, 2009.**

This list is modified from an unpublished list assembled by Manfred Asche. All taxa south of the United States are included. Synonymies may be incomplete, but all published synonymies since Metcalf's (1943) catalog should be included (misspellings noted in the Metcalf catalog are ignored in the synonymy lists). Taxa are listed alphabetically in the current combination. The classification adopted is the most recent available. Most of the distribution records are compiled from published sources (see esp. Muir and Giffard, 1924, and Caldwell and Martorell, 1951), although some are specimen records.

**FAMILY DELPHACIDAE**

<b>Asiracinae</b>	
<b>Asiracini</b> Fieber, 1792	
<i>Asiraca</i> Latreille, 1796 = <i>Manchookhonia</i> Kato, 1933, syn. by Asche 1985: 116.	
<i>germari</i> Metcalf, 1943 = <i>Asiraca longicornis</i> Germar 1830 (nec. <i>Asiraca longicornis</i> Latreille 1802). = <i>Araeopus longicornis</i> (Germar 1830); Schaum, 1850: 69. = <i>Asiraca germari</i> Metcalf, 1943; nom. Nov. for unavailable <i>Asiraca longicornis</i> Germar 1830 (nec. Latreille 1802).	Brazil
<b>Copicerus</b> Swartz, 1802 = <i>Jerala</i> Walker, 1858; syn. by. Stål, 1862: 488. = <i>Holotus</i> Guerin-Meneville, 1856 ; syn. by Stål, 1866: 175.	
<i>insignicornis</i> (Lethierry, 1890) = <i>Asiraca insignicomis</i> Lethierry, 1890. = <i>Copicerus insignicornis</i> (Lethierry, 1890); new comb. by Crawford, 1914: 565.	Venezuela
<i>irroratus</i> Swartz, 1802 = <i>Jerala singula</i> Walker, 1858; syn. by. Stål, 1862: 488.	USA: OH, FL, IL, MS, MO, TX ; Belize, Costa Rica, Cuba; Ecuador, Guatemala, Hispaniola, Jamaica, Mexico, Nicaragua, Panama, Puerto Rico, Venezuela
<i>irroratus thoracicus</i> (Guerin-Meneville, 1856) = <i>Holotus thoracicus</i> Guerin-Meneville, 1866 = <i>Copicerus thoracicus</i> (Guerin-Meneville, 1856); new comb. by Johnson and Fox 1892: 60. = <i>Copicerus irroratus thoracicus</i> (Guerin- Meneville, 1856); new comb. by Crawford, 1914: 564.	
<i>obscurus</i> (Guerin-Meneville, 1856) = <i>Holotus obscurus</i> Guerin-Meneville, 1856.	Brazil

= <i>Copicerus obscurus</i> (Guerin-Meneville, 1856); comb. by Muir, 1926: 7.	
<i>swartzi</i> Stål, 1857	Brazil
<b>Idiosystanini</b> Emeljanov 1996	
<b>Idiosemus</b> Berg, 1883 = <i>Stenosystatus</i> Muir, 1930; syn. by Fennah, 1965: 251.	
<i>xiphias</i> (Berg, 1879) = <i>Liburnia xiphias</i> Berg, 1879. = <i>Idiosemus xiphias</i> (Berg, 1879); new comb. by Berg, 1883. = <i>Stenosystatus anonymi</i> Muir, 1930 ; syn. by Fennah, 1965: 251.	Argentina, Uruguay
<b>Idiosystatus</b> Berg, 1883	
<i>acutiusculus</i> (Spinola, 1852) = <i>Delphax acutiuscula</i> Spinola, 1852. = <i>Idiosystatus acutiusculus</i> (Spinola, 1852); new comb. by Berg, 1883: 231.	Chile, Argentina
<i>australis</i> Fennah, 1969	Argentina
<i>fuscoirroratus</i> (Spinola, 1852) = <i>Delphax fuscoirrorata</i> Spinola, 1852. = <i>Liburnia fuscoirrorata</i> (Spinola, 1852); comb. by Berg, 1881. = <i>Megamelus fuscoirroratus</i> (Spinola, 1852); new comb. by Crawford, 1914: 632. = <i>Delphacodes fuscoirrorata</i> (Spinola, 1852); new comb. by Muir, 1926: 38. = <i>Idiosystatus fuscoirroratus</i> (Spinola, 1852) new comb. by Fennah, 1969: 50-51. = <i>Delphax vittata</i> Spinola, 1852 (nec. Stal, 1862); syn by Fennah, 1969: 50-51. = <i>Delphacodes vittata</i> (Spinola, 1852); new comb. by Muir, 1926: 38.	Chile
<i>gubernator</i> Fennah, 1969	Argentina
<i>longifrons</i> Fennah, 1969	Chile
<b>Pentagramma</b> Van Duzee, 1897 = <i>Bergia</i> Scott, 1881; syn. by Asche, 1985: 283. = <i>Bergias</i> Kirkaldy, 1904; nom. nov. for <i>Bergia</i> Scott, 1881; syn. by Asche, 1985: 283. = <i>Lepticus</i> Crawford, 1914; syn. by Muir, 1926: 3.	
<i>bivittata</i> Crawford, 1914	USA: OH, KS, USA: TX; Nicaragua, British Guiana, Puerto Rico, Trinidad
<i>cosquina</i> Penner, 1947	Argentina, Brazil
<i>nigrifrons</i> Muir, 1934	Argentina
<i>nimbata</i> (Berg, 1879) = <i>Liburnia nimbata</i> Berg 1879 . = <i>Bergia nimbata</i> (Berg, 1879); new comb. by	Argentina

<p>Scott, 1881: 155  = <i>Bergias nimbata</i> (Berg, 1879); new comb.  by Muir, 1926: 36.  = <i>Pentagramma nimbata</i> (Berg, 1879); new  comb by Asche, 1985: 283 by  implication.</p>	
<b>Platysystatini</b> Emeljanov 1996	
<b><i>Equasystatus</i></b> Asche, 1983	
<p><i>breviceps</i> (Muir, 1926)  = <i>Punana breviceps</i> Muir, 1926; new comb.  by Asche, 1983.</p>	Ecuador
<b><i>Platysystatus</i></b> Muir, 1930	
<i>brunneus</i> Muir, 1930	Brazil
<i>itapetingus</i> Asche, 1983	Brazil
<b>Ugyopinae</b>	
<b>Tetrasteirini</b> Emeljanov 1996	
<b><i>Tetrasteira</i></b> Muir, 1926	
<i>albitarsis</i> Fennah, 1945	Trinidad
<i>minuta</i> Muir, 1926	Brazil
<b>Neopunani</b> Emeljanov 1996	
<b><i>Neopunana</i></b> Asche, 1983	
<p><i>alapa</i> (Caldwell, 1951)  = <i>Punana alapa</i> Caldwell, 1951.  = <i>Neopunana alapa</i> (Caldwell, 1951); new  comb. by Asche, 1983: 139.</p>	Puerto Rico
<p><i>caribbensis</i> (Caldwell, 1951)  = <i>Punana caribbensis</i> Caldwell, 1951.  <i>Neopunana caribbensis</i> (Caldwell, 1951); new  comb. by Asche, 1983: 139.</p>	Puerto Rico (Vieques Is.), St. Thomas, Virgin Islands
<p><i>dolon</i> (Fennah, 1959)  = <i>Punana dolon</i> Fennah, 1959  = <i>Neopunana dolon</i> (Fennah, 1959); new  comb. by Asche, 1983: 139.</p>	St. Kitts, Montserrat
<p><i>dominicana</i> (Fennah, 1959)  = <i>Punana dominicana</i> Fennah, 1951.  = <i>Neopunana dominicana</i> (Fennah, 1959);  new comb. by Asche, 1983: 139.</p>	Dominica
<p><i>puertoricensis</i> (Muir, 1918)  = <i>Punana puertoricensis</i> Muir, 1918.  = <i>Neopunana puertoricensis</i> (Muir, 1918);  new comb. by Asche, 1983: 139.</p>	Puerto Rico
<p><i>robusta</i> (Caldwell, 1951)  = <i>Punana robusta</i> Caldwell, 1951;  = <i>Neopunana robusta</i> (Caldwell, 1951); new  comb. by Asche, 1983: 139.</p>	Puerto Rico
<i>saba</i> Asche, 1983	Saba
<p><i>vulgaris</i> (Caldwell, 1951)  = <i>Punana vulgaris</i> Caldwell, 1951.  = <i>Neopunana vulgaris</i> (Caldwell, 1951); new  comb. by Asche, 1983: 139.</p>	Puerto Rico

<b>Ugyopini</b> Fennah, 1979	
<p><i>Ugyops</i> Guerin-Meneville, 1834</p> <ul style="list-style-type: none"> <li>= <i>Bidis</i> Walker, 1857; syn. by Fowler, 1905: 131, 137.</li> <li>= <i>Cona</i> White, 1879; unavailable, replaced by <i>Micromasoria</i> Kirkaldy, 1904.</li> <li>= <i>Consualia</i> Distant, 1917; syn. by Asche, 1985: 283).</li> <li>= <i>Eucanyra</i> Crawford, 1914; syn. by Fennah, 1959: 247.</li> <li>= <i>Epibidis</i> Fowler, 1905; syn. by Fennah, 1959: 247.</li> <li>= <i>Hygiops</i> Amyot and Serville, 1843: 511. syn. by Muir, 1915: 212, 267-269.</li> <li>= <i>Jugodina</i> Schumacher, 1915; syn. by Ishihara, 1949: 9.</li> <li>= <i>Livatis</i> Stål, 1859; syn. by Fennah, 1964: 120; Asche, 1985: 283.</li> <li>= <i>Micromasoria</i> Kirkaldy, 1904; Nom. Nov. for unavailable <i>Cona</i> White, 1879; syn. With <i>Ugyops</i> by Asche, 1985: 283.</li> </ul>	
<p><i>bifurcata</i> (Metcalf, 1945)</p> <ul style="list-style-type: none"> <li>= <i>Eucanyra bifurcata</i>, Metcalf, 1945.</li> <li>= <i>Ugyops bifurcata</i> (Metcalf, 1945); new comb. by Fennah, 1964: 247 (by implication).</li> </ul>	Guyana
<p><i>brunneus</i> (Fowler, 1905)</p> <ul style="list-style-type: none"> <li>= <i>Epibidis brunnea</i> Fowler, 1905.</li> <li>= <i>Ugyops brunneus</i> (Fowler, 1905); new comb. by Fennah, 1964: 123-124.</li> </ul>	Panama
<p><i>flagellata</i> (Fennah, 1945)</p> <ul style="list-style-type: none"> <li>= <i>Eucanyra flagellata</i> Fennah, 1945.</li> <li>= <i>Ugyops flagellata</i> (Fennah, 1945); new comb. by Fennah, 1959: 247.</li> </ul>	Trinidad
<p><i>godmani</i> (Fowler, 1905)</p> <ul style="list-style-type: none"> <li>= <i>Epibidis godmani</i> Fowler, 1905.</li> <li>= <i>Ugyops godmani</i> (Fowler, 1905); new comb. by Fennah, 1964: 122-123.</li> </ul>	?Costa Rica, Panama,
<i>isolata</i> Caldwell, 1951	Puerto Rico (Vieques Is.)
<i>occidentalis</i> Muir, 1918	Puerto Rico
<p><i>osborni</i> Metcalf, 1943</p> <ul style="list-style-type: none"> <li>= <i>Ugyops granulata</i> Osborn, 1935 [nec. <i>Ugyops granulatus</i> Melichar]; unavailable name replaced by Metcalf, 1943: 45.</li> </ul>	Puerto Rico
<p><i>palliatus</i> Fennah, 1964</p> <ul style="list-style-type: none"> <li>= <i>Epibidis godmani</i> Fowler, 1905 (in part); syn. by Fennah, 1964: 121-122.</li> <li>= <i>Epibidis brunnea</i> Fowler, 1905 (in part);</li> </ul>	Panama

syn. by Fennah, 1964: 121-122.	
<i>romani</i> (Muir, 1930) = <i>Eucanyra romani</i> Muir, 1930. = <i>Ugyops romani</i> (Muir, 1930); new comb. by Fennah, 1964: 247 (by implication).	Brazil (Amazons)
<i>stigmata</i> (Crawford, 1914) = <i>Eucanyra stigmata</i> Crawford, 1914. = <i>Ugyops stigmata</i> (Crawford, 1914); new comb. by Fennah, 1964: 247 (by implication).	Belize, Canal Zone
<i>tamu</i> Fennah, 1964	Brazil
<i>taracuae</i> (Muir, 1930) = <i>Eucanyra taracuae</i> Muir, 1930. = <i>Ugyops taracuae</i> (Muir, 1930); new comb. by Fennah, 1964: 247 (by implication).	Brazil (Amazons)
<b><i>Canyra</i> Stål, 1862</b>	
<i>cylindricornis</i> (Fabricius, 1803) = <i>Delphax cylindricornis</i> Fabricius, 1803 = <i>Asiraca cylindricornis</i> (Fabricius, 1803), new comb. by Latreille, 1807: 168. 1830. = <i>Araeopus cylindricornis</i> (Fabricius, 1803) new comb. by Schaum, 1850: 69 . = <i>Canyra cylitidricornis</i> (Fabricius, 1803), new comb. by Stål, 1869: 95.	South America
<i>obscuripennis</i> (Stål, 1854) = <i>Delphax obscuripennis</i> Stål, 1854. = <i>Canyra obscuripennis</i> (Stål, 1854); new comb. by Stål, 1862: 7 (by implication).	Brazil
<i>placida</i> (Stål, 1854) = <i>Delphax placida</i> Stål, 1854. = <i>Canyra placida</i> Stål, 1954); new comb. by Stål, 1862: 7	Brazil
<i>retrahens</i> (Walker, 1858) = <i>Asiraca retrahens</i> Walker, 1858. = <i>Canyra retrahens</i> (Walker, 1858); new comb. by Muir, 1919: 6.	Brazil
<i>revertens</i> (Walker, 1858) = <i>Asiraca revertens</i> Walker, 1858. = <i>Canyra revertens</i> (Walker, 1858); new comb. by Stål, 1862: 488.	Brazil
<i>strigulosa</i> (Walker, 1858) = <i>Asiraca strigulosa</i> Walker, 1858. = <i>Canyra strigulosa</i> (Walker, 1858); new comb. by Stål, 1862: 488.	Brazil
<i>tenensis</i> Muir, 1926	Ecuador
<i>vittifrons</i> (Walker, 1858) = <i>Asiraca vittifrons</i> Walker, 1858	Unknown

= <i>Canyra vittifrons</i> (Walker, 1858); new comb. by Muir, 1919: 6.	
<b>Plesiodelphacinae</b>	
<b>Plesiodelphacini</b> Asche, 1985	
<b><i>Burnilia</i></b> Muir and Giffard, 1924	
<i>belemensis</i> Muir, 1926	?Belize, Brazil, Venezuela
<i>heliconiae</i> Muir, 1926	Guiana
<i>pictifrons</i> (Stål, 1864) = <i>Delphax pictifrons</i> Stål, 1864 = <i>Proterosydne pictifrons</i> (Stål, 1864); new comb. by Crawford, 1914: 571. = <i>Burnilia pictifrons</i> (Stål, 1864); new comb. by Muir and Giffard, 1924: 7.	Belize, Ecuador, Honduras, Mexico, Nicaragua
<i>spinifera</i> Fennah, 1945 = <i>Burnilia spinifera spinifera</i> Fennah, 1945; subspecies by implication Fennah, 1959: 251.	Dominica, Guyana, St. Vincent, Trinidad
<i>spinifera antillana</i> Fennah, 1959	Dominica, St. Vincent
<i>williamsi</i> Muir, 1926	Brazil
<b><i>Plesiodelphax</i></b> Asche, 1985	
<i>guayanus</i> Asche, 1985	Brazil, French Guiana
<i>longicaput</i> (Muir, 1926) = <i>Burnilia longicaput</i> Muir, 1926; new comb. by Asche, 1985.	Brazil, Guyana
<b>Kelisiinae</b>	
<b><i>Kelisiini</i></b> Wagner, 1963	
<b><i>Kelisia</i></b> Fieber, 1866	
<i>vittata</i> Muir, 1926 <sup>1</sup>	Brazil
<b>Stenocraninae</b>	
<b>Stenocranini</b> Wagner, 1963	
<b>Stenocranus</b> Fieber, 1866	
<i>angustus</i> Crawford, 1914	Honduras
<i>maculipes</i> (Berg, 1879)	Argentina
<b>Delphacinae</b>	
<b>Saccharosydniini</b> Asche, 1985	
<b><i>Neomalaxa</i></b> Muir, 1918	
<i>flava</i> Muir, 1918	Brazil, Dominica, Ecuador, Guyana, Puerto Rico, St. Lucia, Trinidad, Venezuela,\
<b><i>Pseudomacrocorupha</i></b> Muir, 1930	
<i>wagneri</i> Muir, 1930	Argentina
<b><i>Saccharosydne</i></b> Kirkaldy, 1907	
<i>brevirostris</i> Muir, 1926	Ecuador
<i>gracillis</i> Muir, 1926	Brazil
<i>ornatipennis</i> Muir, 1926	St. Lucia, Brazil
<i>rostrifrons</i> (Crawford, 1914) = <i>Stenocranus rostrifrons</i> Crawford, 1914. = <i>Saccharosydne rostrifrons</i> (Crawford, 1914);	Caribbean, Cuba, St. Lucia

<sup>1</sup> This species is not a *Kelisia*, it belongs in the Delphacini.

new comb. by Muir and Giffard, 1924: 9.	
<i>saccharivora</i> (Westwood, 1833) = <i>Delphax saccharivora</i> Westwood, 1833. = <i>Stenocranus saccharivora</i> (Westwood, 1822); new comb. by Kirkaldy, 1906: 279, 409. = <i>Saccharosydne saccharivora</i> (Westwood, 1833); new comb. by Kirkaldy, 1907: 139	USA: FL, HA, GA, NC, WI; Barbados, Bermuda, Cuba, Grenada, Haiti, Jamaica, Puerto Rico, St. Lucia, Trinidad, Venezuela.
<i>viridis</i> Muir, 1926	British Guiana
<b>Tropidocephalini</b> Muir, 1915	
<b>Columbiana</b> Muir, 1919	
<i>carasi</i> Fennah, 1964	Peru
<i>lloydi</i> Muir, 1919	Columbia
<b>Columbisoga</b> Muir, 1921	
<i>s.g. Columbisoga</i> Muir, 1921	
<i>chusqueae</i> Muir, 1926	Ecuador
<i>filistylus</i> Muir, 1930	Bolivia
<i>gynerii</i> Muir, 1926	Ecuador
<i>gyneriicola</i> Muir, 1926	Ecuador
<i>maculosa</i> Muir, 1926	Ecuador
<i>merae</i> Muir, 1926	Ecuador
<i>ornata</i> Muir, 1926	Ecuador
<i>sacchari</i> Muir, 1926	Ecuador
<i>saracura</i> Fennah, 1963	Brazil
<i>tenae</i> Muir, 1926	Ecuador
<i>zapote</i> Muir, 1926	Ecuador
<b>s.g. Columbisodes</b> Fennah, 1963	
<i>tylotus</i> Fennah, 1964	Brazil
<b>Macrocorupha</b> Muir, 1926	
<i>gynerii</i> Muir, 1926	Ecuador
<b>Malaxa</b> Melichar, 1914	
<i>gracilis</i> Fennah, 1945	Venezuela
<i>microstyla</i> Muir, 1930	Bolivia
<i>occidentalis</i> Muir, 1926	Ecuador
<b>Delphacini</b> Leach, 1815	
<b>Abrosoga</b> Caldwell, 1951	
<i>errata</i> Caldwell, 1951	Puerto Rico
<b>Anchidelphax</b> Fennah, 1965	
<i>albidens</i> (Crawford, 1914) = <i>Megamelus albidens</i> Crawford, 1914 = <i>Delphacodes albidens</i> (Crawford, 1914); new comb. by Muir and Giffard, 1924: 28. = <i>Anchidelphax albidens</i> (Crawford, 1914); new comb. by Fennah, 1965: 97.	Brazil
<i>havanensis</i> (Crawford, 1914) = <i>Megamelus albidens havanensis</i> Crawford,	Cuba, Jamaica, Martinique, Puerto Rico, St. Lucia

1914. = <i>Delphacodes havanensis</i> (Crawford); new comb. by Fennah, 1965: 97. = <i>Delphacodes mardininae</i> Muir, 1918; syn. by Muir and Giffard, 1924: 28.	
<b>Caenodelphax</b> Fennah, 1965	
<i>philyra</i> (Fennah, 1959) = <i>Delphacodes philyra</i> Fennah, 1959. = <i>Caenodelphax philyra</i> (Fennah, 1969); new comb. by Fennah, 1965: 96.	St. Lucia
<i>teapae</i> (Fowler, 1905) = <i>Liburnia teapae</i> Fowler, 1905. = <i>Delphacodes teapae</i> (Fowler, 1905); new comb. by Wolcot, 1923; 274. = <i>Caenodelphax teapae</i> (Fowler, 1905); new comb. by Fennah, 1965: 96 (from <i>Delphacodes</i> ).	USA: FL; Barbados, Belize, Brazil, Costa Rica, Cuba, Dominica, Ecuador, Guyana, Guadeloupe, Jamaica, Martinique, Mexico, Montserrat, Nicaragua, Puerto Rico, St. Lucia, St. Vincent, Trinidad
<b>Calbodus</b> Spinola, 1852	
<i>pallidulus</i> Spinola, 1852 = <i>Delphacodes correntosoensis</i> Muir, 1929; syn. by Fennah, 1965: 251.	Chile, Argentina
<i>patquianus</i> Fennah, 1965	Argentina
<b>Chionomus</b> Fennah, 1971	
<i>balboae</i> (Muir and Giffard, 1924) = <i>Delphacodes balboae</i> Muir and Giffard, 1924. = <i>Chionomus balboae</i> (Muir and Giffard, 1924); new comb. by Fennah, 1971: 324.	USA: FL; Argentina, Mexico, Ecuador, Guyana, Mexico, Puerto Rico, St. Lucia, Venezuela
<i>havanae</i> (Muir and Giffard, 1924) = <i>Delphacodes havanae</i> Muir & Giffard, 1924. = <i>Chionomus havanae</i> (Muir and Giffard, 1924); new comb. by Fennah, 1971: 324.	USA: FL; Cayman Is. (Grand Cayman), Cuba, Jamaica, Guatemala, Guyana, Puerto Rico, Trinidad, Venezuela
<i>haywardi</i> (Muir, 1929) = <i>Delphacodes haywardi</i> Muir, 1929. = <i>Chionomus haywardi</i> (Muir, 1929); new comb. by Fennah, 1971: 324. = <i>Delphacodes collaris</i> De Remes, Lenikov & Teson, 1978; syn. by Teson and Remes Lenicov, 1983: 322.	Argentina, Uruguay
<b>Chloriona</b> Fieber, 1866	
<i>nigrifrons</i> Van Duzee, 1908	Jamaica
<b>Delphacodes</b> Fieber, 1866	
<i>aculeata</i> Beamer, 1948	USA: FL
<i>andromeda</i> (Van Duzee, 1907) = <i>Liburnia andromeda</i> Van Duzee, 1907. = <i>Delphacodes andromeda</i> (Van Duzee,	USA: FL, IL, MA, MO, MS, NC, TN; Cuba, Guyana, Honduras, Jamaica, Puerto Rico,

1907); new comb. by Muir and Giffard, 1924: 36. = <i>Pissonotus speciosus</i> Metcalf, 1923; syn. by Morgan and Beamer, 1949: 142.	
<i>ardentis</i> Beamer, 1948	USA: FL; Mexico
<i>argentina</i> De Remes, Lenikov & Teson, 1979	Argentina
<i>arcuata</i> beamer, 1948 = <i>Delphacodes albinotata</i> Muir and Giffard, 1924 (nec. Crawford, 1914); replacement name by Beamer, 1948: 106. = <i>Delphacodes arcuata</i> Beamer, 1948; replacement name for unavailable <i>D. albinotata</i> Muir and Giffard, 1924.	Dominica, Guadeloupe, Mexico, Puerto Rico (inc. Vieques Is.)
<i>aterrima</i> Muir, 1926	Brazil, Guyana, ?Surinam
<i>atrior</i> (Fowler, 1905) = <i>Liburnia atrior</i> Fowler, 1905. = <i>Megamelus atrior</i> (Fowler, 1905); new comb. by Crawford, 1914: 608, 628, 630. = <i>Delphacodes atrior</i> (Fowler, 1905); new comb. by Metcalf, 1943: 407.	Mexico
<i>banosensis</i> Muir, 1926	Ecuador
<i>bergi</i> (Scott, 1881) = <i>Liburnia cognate</i> Berg, 1879 [nec. Fieber, 1866]. = <i>Liburnia bergi</i> Scott, 1881: 156. nom. nov. for preoccupied <i>Liburnia cognata</i> Berg [nec Fieber]. = <i>Megamelus bergi</i> (Scott, 1881); new comb. by Crawford, 1914 :631. = <i>Delphacodes bergi</i> (Scott, 1881); new comb. by Muir, 1926: 38.	Argentina
<i>caymanensis</i> (Crawford, 1914) = <i>Megamelus cayamensis</i> Crawford, 1914 = <i>Delphacodes cayamensis</i> (Crawford, 1914); new comb. by Muir, and Giffard, 1924: 29.	Cuba
<i>chiloensis</i> Muir, 1929	Chiloe Is. (Chile)
<i>culta</i> (Van Duzee, 1907) = <i>Liburnia culta</i> Van Duzee, 1907. = <i>Megamelus cultus</i> (Van Duzee, 1907); new comb. by Crawford, 1914 :628. = <i>Delphacodes culta</i> (Van Duzee, 1907); new comb. by Muir and Giffard, 1924: 38.	USA: FL: Bermuda, Cuba, Jamaica,
<i>darwini</i> Muir, 1929	Chiloe Is. (Chile)
<i>detecta</i> (Van Duzee, 1897) = <i>Liburnia detecta</i> Van Duzee, 1897 = <i>Liburnia circumcincta</i> Van Duzee, 1909; syn. by Van Duzee, 1912: 506.	USA: CT, GA, FL, LA, MA, MD, MS, NC, NJ, NY, TX, VA; Bermuda, Puerto Rico (inc. Vieques Is.), St. Thomas

<p>= <i>Megamelus circumcinctus</i> (Van Duzee, 1909); new comb. by Crawford, 1914.  = <i>Megamelus vanduzeei</i> Crawford, 1914; syn. by Muir and Giffard, 1924: 26.  = <i>Liburnia vanduzeei</i> (Crawford, 1914); new comb. by Van Duzee, 1916, 84.  = <i>Delphacodes detecta</i> (Van Duzee, 1897); comb. by Muir and Giffard, 1924: 26.</p>	
<i>elongatus</i> Teson & Remes Lenicov, 1983	Argentina
<i>floridae</i> Muir and Giffard, 1924	USA:FL, Jamaica, Montserrat, Puerto Rico,
<i>floridae puertoricensis</i> Caldwell, 1951	Puerto Rico, Montserrat
<p><i>fulvidorsum</i> (Metcalf, 1923)  = <i>Liburnia fulvidorsum</i> Metcalf, 1923.  = <i>Delphacodes fulvidorsum</i> (Metcalf, 1923); new comb. by Frison, 1927: 153.</p>	USA: FL, GA,MO, TX, MO; Bermuda, Cuba, Guadeloupe, Martinique, Puerto Rico
<p><i>fuscoterminata</i> (Berg, 1879)  = <i>Liburnia fuscoterminata</i> Berg, 1879.  = <i>Megamelus fuscoterminata</i> (Berg, 1879); new comb. by Crawford, 1914: 632.  = <i>Delphacodes fuscoterminata</i> (Berg, 1879); new comb. by Muir, 1926: 38.</p>	Argentina
<i>gluciophila</i> Muir, 1926	Ecuador
<i>guianensis</i> Muir, 1919	Guiana
<p><i>humilis</i> (Van Duzee, 1907)  = <i>Liburnia humilis</i> Van Duzee, 1907.  = <i>Megamelus humilis</i> (sic) (Van Duzee, 1907); new comb. by Crawford, 1914: 608, 629.  = <i>Megamelus erectus</i> Crawford, 1914; syn. by Muir and Giffard, 1924: 32.  = <i>Delphacodes erectus</i> (Crawford, 1914); new comb. by Muir, 1918: 427.  = <i>Delphacodes humilis</i> (Van Duzee, 1907); new comb. by Wolcott, 1923: 274.</p>	USA: FL; Argentina, Brazil, Cuba, Ecuador, Guadalupe, Guyana, Hispaniola, Jamaica, Mexico, Montserrat, Nicaragua, Puerto Rico, Venezuela
<i>koebeleri</i> Muir & Giffard, 1924	Mexico, Ecuador
<i>kuscheli</i> Fennah, 1955	Chile, Juan Fernandez Is., Argentina
<p><i>latus</i> De Remes, Lenikov &amp; Teson, 1979  = <i>Delphacodes lata</i> De Remes, Lenikov &amp; Teson, 1979; emendation by Remes Lenicov &amp; Virla, 1999: 12.</p>	Argentina
<p><i>lutulenta</i> (Van Duzee, 1897)  = <i>Liburnia lutulenta</i> Van Duzee, 1897.  = <i>Megamelus lutulentus</i> (Van Duzee, 1897); new comb. by Crawford, 1914: 606, 617.  = <i>Delphacodes lutulenta</i> (Van Duzee, 1897); new comb. by Muir and Giffard, 1924: 23.</p>	USA: AZ, DE, CA, CO, CT, KA, KS, KY, IL, IN, MA, MD, ME, MN, MO, NC, NH, NY, OH, PA, SD, TN, VA, VT, WV; CAN: Alberta, British Columbia, Nova Scotia, Quebec, Ontario, Saskatchewan; Puerto Rico
<p><i>marginicornis</i> (Fowler, 1905)  = <i>Liburnia marginicomis</i> Fowler, 1905.</p>	Mexico

<p>= <i>Megamelus marginicomis</i> Crawford, 1914 : 630.  = <i>Delphacodes marginicornis</i> (Fowler, 1905); new comb. by Metcalf, 1943: 448.</p>	
<i>mesada</i> Caldwell 1951	Puerto Rico
<i>montezumae</i> Muir & Giffard, 1924	Mexico
<i>nigrifacies</i> Muir, 1918	USA: FL; Costa Rica, Dominica, Guyana, Martinique, Montserrat, St. Lucia, St. Vincent
<i>nigrinota</i> Beamer, 1951	Mexico
<p><i>nigripennis</i> (Crawford, 1914)  = <i>Megamelus erectus nigripennis</i> Crawford, 1914.  = <i>Delphacodes erectus nigripennis</i> (Crawford, 1914); new comb. by Muir, 1918: 427.  = <i>Delphacodes nigripennis</i> (Crawford, 1914); new comb. by Muir and Giffard, 1924: 31.</p>	Cuba, Jamaica, Martinique, Nicaragua, Puerto Rico,
<p><i>pacifica</i> (Crawford, 1914)  = <i>Megamelus pacificus</i> Crawford, 1914.  = <i>Liburnia pacifica</i> (Crawford, 1914); new comb. by Van Duzee, 1916: 84.  = <i>Delphacodes pacifica</i> (Crawford, 1914); new comb. by Muir and Giffard, 1924: 34.</p>	USA: AZ, CA; Mexico
<i>paludicola</i> Muir, 1926	Ecuador
<p><i>patruelis</i> (Stål, 1859)  = <i>Delphax patruelis</i> Stal, 1859.  = <i>Liburnia patruelis</i> (Stål, 1859); new comb. by Berg, 1879: 188.  = <i>Megamelus patruelis</i> (Stål, 1859); new comb. by Crawford, 1914 :632.  = <i>Delphacodes patruelis</i> (Stål, 1859); new comb. by Muir, 1926: 37.</p>	Argentina
<i>platystylus</i> Muir, 1930	Argentina
<p><i>puella</i> (Van Duzee, 1897)  = <i>Liburnia puella</i> Van Duzee, 1897.  = <i>Delphax puella</i> (Van Duzee, 1897); new comb. by Kirkaldy, 1907: 161.  = <i>Megamelus puella</i> (Van Duzee, 1897); new comb. by Crawford, 1914: 622, 625-626.  = <i>Megamelus puellis</i> (Van Duzee, 1897); missp. By Crawford, 1914: 625-626.  = <i>Delphacodes puella</i> (Van Duzee, 1897); new comb. by Muir and Giffard, 1924: 32.</p>	USA: AL, AR, CA, CT, CO, DC, FL, GA, HI, IA, IL, IN, KS, KY, LA, MA, MD, ME, MO, MS, NC, NE, NH, NJ, NY, OH, OK, OR, PA, TN, VA, WA; Canada: Que., Ont.; Bermuda, Brazil, Cuba, Mexico, Puerto Rico,
<i>quadrispinosa</i> Muir & Giffard, 1924	Cuba, Nicaragua
<p><i>rectangularis</i> (Crawford, 1914)  = <i>Megamelus rectangularis</i> Crawford, 1914.</p>	Nicaragua

= <i>Delphacodes rectangularis</i> (Crawford, 1914); new comb. by Metcalf, 1943: 502.	
<i>reducta</i> (Van Duzee, 1907) = <i>Liburnia</i> (?) <i>reducta</i> Van Duzee, 1907. = <i>Megamelus reductus</i> (Van Duzee, 1907); Crawford, 1914: 629. = <i>Delphacodes reducta</i> (Van Duzee, 1907); new comb. by Metcalf, 1943: 502.	Jamaica
<i>saccharicola</i> Muir, 1926	Ecuador
<i>sagata</i> (Fowler, 1905) = <i>Liburnia sagata</i> Fowler, 1905. = <i>Megamelus sagatus</i> (Fowler, 1905); Crawford, 1914: 622, 630. = <i>Delphacodes sagata</i> (Fowler, 1905); new comb. by Metcalf, 1938: 301.	Canal Zone, Mexico
<i>saxicola</i> Muir, 1926	Brazil
<i>securigera</i> Muir, 1926	Brazil
<i>seminigra</i> (Stål, 1854) = <i>Delphax seminigra</i> Stål, 1854. = <i>Liburnia seminigra</i> (Stål, 1854); new comb. by Van Duzee, 1907:46. = <i>Megamelus seminigra</i> (Stål, 1854); new comb. by Crawford, 1914, 631; Barber, 1914: 528. = <i>Delphacodes seminigra</i> (Stål, 1854); new comb. by Muir and Giffard, 1924: 38.	USA: CO, FL, Brazil, Jamaica
<i>sitarea</i> De Remes, Lenikov & Teson, 1979 = <i>Delphacodes sitare</i> , missp. By De Remes, Lenikov & Teson, 1979: 75.	Argentina
<i>spinus</i> Teson and Remes Lenicov, 1983	Argentina
<i>tenae</i> Muir, 1926	Ecuador
<i>univittata</i> (Berg, 1879) = <i>Liburnia univittata</i> Berg, 1879. = <i>Megamelus univittata</i> (Berg, 1879); new comb. by Crawford, 1914: 632. = <i>Delphacodes univittata</i> (Berg, 1879); new comb. by Muir, 1926.	Argentina
<i>vaccina</i> Caldwell, 1951	Puerto Rico
<i>xerophila</i> Muir, 1926	Ecuador
<i>xerosa</i> Caldwell, 1951	Puerto Rico
<b><i>Delphax</i></b> Fabricius, 1798 = <i>Araeopus</i> Spinola, 1839 (ICZN, 1961)	
<i>setosus</i> (Germar, 1830) = <i>Aeropus setosus</i> (Germar, 1830); <i>Delphax</i> senior synonym of <i>Aeropus</i> (ICZN, 1961). = <i>Asiraca setosus</i> Germar, 1830; originally to <i>Aeropus</i> by Schaum, 1850.	Brazil
<i>conspersinervis</i> (Lethierry, 1890)	Venezuela

<p>= <i>Araeopus conspersinervis</i> Lethierry, 1890.  = <i>Delphax conspersinervis</i> (Lethierry, 1890);  new comb. by Crawford, 1914: 577.</p>	
<p><i>producta</i> (Walker, 1851)  = <i>Delphax producta</i> Walker, 1851.  = <i>Araeopus productus</i> Walker, 1851; new  comb. by Metcalf, 1943: 384.  Restored to <i>Delphax</i> by ICZN, 1961.</p>	Jamaica
<b><i>Dicranotropis</i></b> Distant, 1906	
<i>acheron</i> Fennah, 1957	Chile: Juan Fernandez Island
<p><i>basifuscata</i> (Fowler, 1905)  = <i>Liburnia basifuscata</i> Fowler, 1905  = <i>Megamelus basifuscatus</i> (Fowler, 1905);  comb. by Crawford, 1914: 608. 630.  = <i>Dicranotropis basifuscata</i> (Fowler, 1905);  comb. by Remes Lenicov, 1996: 124</p>	Argentina, Mexico
<i>bipectinata</i> Muir, 1926	Ecuador
<p><i>fuscoterminata</i> (Berg, 1879)  = <i>Liburnia fuscoterminata</i> Berg, 1879.  = <i>Megamelus fuscoterminata</i> (Berg, 1879);  new comb. by Crawford, 1914: 632.  = <i>Dicranotropis fuscoterminata</i> (Berg, 1879);  syn. by Remes Lenicov, 1996: 124.  = <i>Liburnia nigricula</i> Berg, 1879; syn. by  Remes Lenicov, 1996: 124.  = <i>Megamelus nigricula</i> (Berg, 1879); comb.  by Crawford, 1914: 632.  = <i>Delphacodes nigricula</i> (Berg, 1879); comb.  by Muir, 1926: 37.</p>	Argentina
<b><i>Euidelloides</i></b> Muir, 1926	
<i>montana</i> Muir, 1926	Ecuador
<p><b><i>Euides</i></b> Fieber, 1866  = <b><i>Euidella</i></b> Fieber, 1866; syn. by Metcalf, 1952:  230-231.</p>	
<p><i>afasciata</i> (Caldwell, 1951)  = <i>Euidella afasciata</i> Caldwell, 1951.  = <i>Euides afasciata</i> (Caldwell, 1951); comb.  by Metcalf, 1952: 230-231 (by  implication).</p>	Puerto Rico, St. Lucia
<p><i>altamazonica</i> (Muir, 1926)  = <i>Euidella altamazonica</i> Muir, 1926.  = <i>Euides altamazonica</i> (Muir, 1926); comb.  by Metcalf, 1952: 230-231 (by  implication).</p>	Ecuador
<p><i>belemensis</i> (Muir, 1926)  = <i>Pissonotus belemensis</i> Muir, 1926.  = <i>Euides belemensis</i> (Muir, 1926); new comb.  by Bartlett &amp; Deitz, 2000: 146.</p>	Brazil
<p><i>blairmontensis</i> (Muir, 1926)  = <i>Euidella blairmontensis</i> Muir, 1926.</p>	Guiana

<p>= <i>Euides blairmontensis</i> (Muir, 1926); comb. by Metcalf, 1952: 230-231 (by implication).</p>	
<p><i>brazilensis</i> (Muir, 1926)  = <i>Pissonotus brazilensis</i> Muir, 1926.  = <i>Euides brazilensis</i> (Muir, 1926); new comb. by Bartlett &amp; Deitz, 2000: 146.</p>	Brazil, Ecuador, St. Lucia
<p><i>elegans</i> (Muir, 1926)  = <i>Euidella elegans</i> Muir, 1926.  = <i>Euides elegans</i> (Muir, 1926); comb. by Metcalf, 1952: 230-231 (by implication).</p>	Ecuador
<p><i>equadorensis</i> (Muir, 1926)  = <i>Pissonotus equadorensis</i> Muir, 1926.  = <i>Euides equadorensis</i> (Muir, 1926); new comb. by Bartlett &amp; Deitz, 2000: 146.</p>	Ecuador
<p><i>fasciatella</i> (Osborn, 1935)  = <i>Liburniella fasciatella</i> Osborn, 1935.  = <i>Euidella fasciatella</i> (Osborn, 1935); new comb. by Caldwell and Martorell, 1951: 190.  = <i>Euides fasciatella</i> (Osborn, 1935); comb. by Metcalf, 1952: 230-231 (by implication).  = <i>Pissonotus striolus</i> Osborn, 1935; syn. by Caldwell and Martorell, 1951: 190.</p>	Puerto Rico
<p><i>fucata</i> Berg, 1883  = <i>Euides fucata</i> Berg, 1883; comb. restored by Metcalf, 1952: 230-231 (by implication).  = <i>Megamelus fucatus</i> (Berg, 1883); new comb. by Crawford, 1914: 632.  = <i>Euidella fucata</i> (Berg, 1883); comb. by Muir, 1926: 37.</p>	Argentina
<p><i>fuscovittata</i> (Scott, 1881)  = <i>Euides fuscovittata</i> Scott, 1881; comb. restored by Metcalf, 1952: 230-231 (by implication).  = <i>Megamelus fusco-vittata</i> (Scott, 1881); new comb. by Crawford, 1914: 632.  = <i>Euidella fusco-vittata</i> (Scott, 1881); new comb. by Muir, 1926: 37.</p>	Argentina
<p><i>grossa</i> (Van Duzee, 1933)  = <i>Euidella grossa</i> Van Duzee, 1933.  = <i>Euides grossa</i> (Van Duzee, 1933); comb. by Metcalf, 1952: 230-231 (by implication).</p>	Costa Rica
<p><i>guaduae</i> (Muir, 1926)  = <i>Euidella guaduae</i> Muir, 1926.  = <i>Euides guaduae</i> (Muir, 1926); comb. by</p>	Ecuador

Metcalf, 1952: 230-231 (by implication).	
<i>magnistyla</i> (Crawford, 1914) = <i>Megamelus magnistylus</i> Crawford, 1914. = <i>Euidella magnistylus</i> (Crawford, 1914); new comb. by Muir and Giffard, 1924: 10. = <i>Euides magnistylus</i> (Crawford, 1914); comb. by Metcalf, 1952: 230-231 (by implication).	USA: GA, Cuba, Mexico, Puerto Rico
<i>megalostylus</i> (Muir, 1919) = <i>Pissonotus megalostylus</i> Muir, 1919 = <i>Euides megalostylus</i> (Muir, 1919); new comb. by Bartlett & Deitz, 2000: 146.	Guiana
<i>triloba</i> (Metcalf, 1923) = <i>Liburnia triloba</i> Metcalf, 1923. = <i>Euidella triloba</i> (Metcalf, 1923). = <i>Euides triloba</i> (Metcalf, 1923); comb. nov. by Metcalf, 1952: 230-231 (by implication).	USA: FL, LA, MS; Bermuda
<i>weedi</i> (Van Duzee, 1897) = <i>Liburnia weedi</i> Van Duzee, 1897. = <i>Euidella weedi</i> (Van Duzee, 1897); new comb. by Muir and Giffard, 1924: 10. = <i>Euides weedi</i> (Crawford, 1914); comb. by Metcalf, 1952: 230-231 (by implication).	USA: FL, GA, IL, KY, MO, MS, TX; Bermuda, Cuba, Mexico, Puerto Rico.
<b><i>Eurysa</i></b> Fieber, 1866	
<i>dorsilinea</i> (Van Duzee, 1907) = <i>Liburnia</i> (?) <i>dorsilinea</i> Van Duzee, 1907. = <i>Megamelus dorsilinia</i> [sic] (Van Duzee, 1907); comb by Crawford, 1914: 631. = <i>Eurysa dorsilinia</i> [sic] (Van Duzee, 1907); comb. by Muir and Giffard, 1924: 9.	Cuba, Jamaica
<b><i>Falcotoya</i></b> Fennah, 1969	
<i>crawfordi</i> (Muir & Giffard, 1924) = <i>Delphacodes crawfordi</i> Muir and Giffard, 1924. = <i>Falcotoya crawfordi</i> (Muir and Giffard, 1924); comb. by Fennah, 1969: 21.	Mexico
<b><i>Hadropygos</i></b> Gonzon and Bartlett, 2008	
<i>rhombos</i> Gonzon and Bartlett, 2008	Bolivia
<b><i>Javesella</i></b> Fennah, 1963 = <i>Weidnerianella</i> Wagner, 1963; syn. by Le Quesne, 1964: 57.	
<i>pellucida</i> (Fabricius, 1794) = <i>Fulgora pellucida</i> Fabricius, 1794. = <i>Fulgora striata</i> Fabricius, 1794; syn. with <i>Delphax dispar</i> Fallén, 1806 by Zetterstedt, 1828: 548; syn. with <i>Delphax pellucida</i> by Flor, 1861: 65.	Widely distributed in North temperate regions.  USA: AK, CO, CT, IL, MA, ME, MI, NC, ND, NH, NJ, NV, NY, OH, OR, PA, WA, WI, WY; CAN: Alberta, British Columbia, Ontario, Quebec); Cuba, Puerto Rico.

- = *Fulgora marginata* Fabricius, 1794; syn. with *Delphax dispar* Fallén, 1806 by Fallén, 1806: 126; syn. with *Delphax pellucida* by Herrich-Schäffer, 1836: 18.
- = *Fulgora flavescens* Fabricius, 1794; syn. with *Delphax dispar* Fallén, 1806 by Fallén, 1806: 126; syn. with *Delphax pellucida* by Flor, 1861: 65.
- = *Cicada pellucida* (Fabricius, 1794); new comb. by Weber, 1795: 146.
- = *Delphax pellucida* (Fabricius, 1794); new comb. by Fabricius, 1803 :84.
- = *Delphax striata* (Fabricius, 1794); new comb. by Fabricius, 1803: 84.
- = *Delphax marginata* (Fabricius, 1794); new comb. by Fabricius, 1803: 84.
- = *Delphax flauescens* [sic] (Fabricius, 1794); new comb. by Fabricius, 1803: 84.
- = *Asiraca pellucida* (Fabricius, 1794); new comb. by Latreille, 1804: 318.
- = *Asiraca striata* (Fabricius, 1794); new comb. by Latreille, 1804: 318.
- = *Asiraca marginata* (Fabricius, 1794); new comb. by Latreille, 1804: 318.
- = *Asiraca flavescens* (Fabricius, 1794); new comb. by Latreille, 1804: 318.
- = *Delphax dispar* Fallén, 1806; syn. with *Delphax pellucida* by Flor, 1861: 65..
- = *Delphax suturalis* Curtis, 1837; syn. with *Delphax pellucida* by Flor, 1861: 65.
- = *Delphax dubia* Kirschbaum, 1868 (nec. Panzer, 1796); syn. with *Liburnia pellucida* (Fabricius, 1794) by Fieber, 1872: 5.
- = *Delphax piceola* Kirschbaum, 1868; syn. with *Delphax pellucida* (Fabricius, 1794) by Puton, 1886: 73.
- = *Liburnia pellucida* (Fabricius, 1794); new comb. by Stål, 1869: 117.
- = *Liburnia striata* (Fabricius, 1794); new comb. by Stål, 1869: 117.
- = *Liburnia marginata* (Fabricius, 1794); new comb. by Stål, 1869: 117.
- = *Liburnia flavescens* (Fabricius, 1794); new comb. by Stål, 1869: 117.
- = *Achorotile pellucida* (Fabricius, 1794); new comb. by Oshanin, 1870: 48.
- = *Liburnia suturalis* (Curtis, 1837); new comb. by Fieber, 1872: 5.
- = *Delphax pellucida dispar* (Fallén, 1806);

Algeria, Austria, Belgium, British Isles (inc. Wales), Czechoslovakia, Denmark, Estonia, Finland (Lapland), France, Germany (inc. Baltrum Island), Hungary, Iceland, Ireland, Italy, Japan, Libya, Morocco, Netherlands (inc. Nijmegen as “Batavia”), Norway, Poland, Romania, Russia (Caucasus, Kamchatka, Sakhalin Island, Siberia, Transcaucasia), Sardinia, Scandinavia, Scotland, Spain, Sweden, Switzerland, Turkestan, Yugoslavia.

<p>new comb. by Mayr, 1884 :20.  = <i>Liburnia pellucida dispar</i> (Fallén, 1806);  new comb. by Melichar, 1896: 71.  = <i>Liburnia yezoana</i> Matsumura, 1900; syn. by  Muir and Giffard, 1924: 20.  = <i>Embolophora pellucida</i> (Fabricius, 1794);  new comb. by Kirkaldy, 1901: 338.  = <i>Delphax yezoana</i> (Matsumura, 1900); new  comb by Oshanin, 1907: 330.  = <i>Megamelus pellucidus</i> (Fabricius, 1794);  new comb. by Crawford, 1914: 607,  615.  = <i>Liburnia semicinctus</i> Van Duzee, 1917; syn.  with <i>Delphacodes pellucida</i> by Muir  and Giffard, 1924: 20.  = <i>Delphacodes pellucida</i> (Fabricius, 1794);  new comb. by Muir and Giffard,  1924: 20.  = <i>Javesella pellucida</i> (Fabricius, 1794); new  comb. by Fennah, 1963: 15.</p>	
<b>Kormus</b> Fieber, 1866	
<i>apicimaculatusa</i> (Fowler, 1905)	Guatemala
<b>Liburnia</b> Stål, 1866	
<p><i>mexicana</i> (Crawford, 1914)  = <i>Megamelus puellis mexicanus</i> Crawford,  1914.  = <i>Sogata mexicana</i> (Crawford, 1914); new  comb by Muir and Giffard, 1924: 15  = <i>Liburnia mexicana</i> (Crawford, 1914); comb.  by Metcalf, 1943: 365.</p>	Mexico
<p><i>paludata</i> Fowler, 1905  = <i>Sogata paludata</i> Fowler; new comb by.  Muir and Giffard, 1924.  = <i>Liburnia paludata</i> Fowler, 1905; comb.  restored by Metcalf, 1943: 366.</p>	Guatemala, Mexico
<p><i>williamsi</i> (Muir, 1926)  = <i>Sogata williamsi</i> Muir, 1926.  = <i>Liburnia williamsi</i> (Muir, 1926); new comb  by. Metcalf, 1943: 371.</p>	Ecuador
<b>Liburniella</b> Crawford, 1914	
<p><i>ornata</i> (Stål, 1862)  = <i>Delphax ornata</i> Stål, 1862.  = <i>Liburnia ornata</i> (Stål, 1862); new comb. by  Osborn, 1892: 127.  = <i>Liburniella ornata</i> (Stål, 1862); new comb.  by Crawford, 1914: 586, 601.</p>	US: AL, CA, CT, DC, DE, FL, GA, IA, IL, IN, KS, KY, LA, MA, ME, MO, MS, NB, NC, NJ, NY, OH, PA, SC, TN, VA; Bermuda [Ecuador error]
<b>Macrotomella</b> Van Duzee, 1907	
<i>carinata</i> Van Duzee, 1907	Jamaica
<b>Matutinus</b> Distant, 1917	

<i>cubensis</i> Rodriguez-Leon et al., 1998	Cuba
<i>fuscipennis</i> (Muir, 1919) = <i>Chloriona fuscipennis</i> Muir, 1919. = <i>Matutinus fuscipennis</i> (Muir, 1919); new comb. by Fennah, 1963: 48.	Brazil
<b>Megamelus</b> Fieber, 1866	
<i>bellicus</i> Remes Lenicov & Sosa, 2007	Argentina, Brazil, Peru
<i>bifurcatus</i> Crawford, 1914	Brazil
<i>electrae</i> Muir, 1926	Brazil, Puerto Rico
<i>iphigeniae</i> Muir, 1926	Brazil
<i>scutellaris</i> Berg, 1883	Argentina
<i>timehri</i> Muir, 1919	Guiana
<b>Metadelphax</b> Wagner, 1963 = <i>Metadelphax</i> Wagner, 1963. Syn. with <i>Toya</i> by Fennah 1964: 142 (by implication; type species moved to <i>Toya</i> ); Nast 1972:65, Linnavuori 1973: 107 (generic synonymy listed). Name restored by Ding 2006: 511. = <i>Meadelphax</i> Wagner, 1963: Missp. by Kuoh et al. 1983: 155.	
<i>argentinensis</i> (Muir, 1929) = <i>Delphacodes variabilis</i> De Remes, Lenikov & Teson, 1978; syn. by Teson & Remes Lenicov 1983: 322. = <i>Delphacodes argentinensis</i> Muir, 1929 = <i>Syndelphax argentinensis</i> (Muir, 1929); comb. by Fennah, 1971: 323. = <i>Toya argentinensis</i> (Muir, 1929); new comb. by Teson & Remes Lenicov 1989: 101-102 = <i>Metadelphax argentinensis</i> (Muir, 1929); New comb. by Gonzon & Bartlett 2008: 234-237.	Argentina, Bolivia, Columbia, Paraguay, Peru, Venezuela
<i>dentata</i> Gonzon and Bartlett, 2008	Ecuador
<i>pero</i> (Fennah, 1971) = <i>Syndelphax pero</i> Fennah, 1971; new comb. by Gonzon and Bartlett, 2008.	Cayman Is., (Grand Cayman); Cuba
<i>propinqua</i> (Fieber, 1866) = <i>Delphax propinqua</i> Fieber, 1866: 525. = <i>Delphax hamulata</i> Kirschbaum, 1868: 38: synonymy by Fieber, 1872: 5. = <i>Liburnia propinqua</i> (Fieber): new combination by Fieber, 1872: 5. = <i>Liburnia marshalli</i> Scott, 1873: 104: synonymy by Webb and Wilson, 1986: 286. = <i>Liburnia terminalis</i> Van Duzee, 1907: 49: synonymy by Muir and Giffard, 1924: 31.	Widely distributed in Tropical and warm temperate regions.  NEW WORLD: USA AZ, AR, CA, DE, FL, GA, IL, KS, KY, LA, MD, MO, MS, NC, OH, SC, TN, TX, UT, WA; Antigua, Argentina, Bahamas, Barbados, Belize, Bermuda, Bolivia, Brazil, CANADA: British Columbia; Cayman Islands, Chile, Colombia, Costa Rica, Cuba, Dominica, Ecuador, El Salvador, Granada, Guatemala, Guyana, Hispaniola, Honduras, Jamaica, Martinique, Mexico, Montserrat, Nicaragua,

<ul style="list-style-type: none"> <li>= <i>Delphax cataniae</i> Matsumura, 1910: 35: synonymy by Nast, 1975: 4.</li> <li>= <i>Delphax graminicola</i> Matsumura, 1910: 17: synonymy by Yang, 1989: 4.</li> <li>= <i>Liburnia tuckeri</i> Van Duzee, 1912: 506: synonymy by Muir and Giffard, 1924: 31.</li> <li>= <i>Delphacodes propinqua</i> (Fieber): new combination by Muir, 1917: 335.</li> <li>= <i>Delphacodes neopropinqua</i> Muir, 1917: 335: synonymy by Muir and Giffard, 1924: 31.</li> <li>= <i>Delphacodes subfusca</i> Muir, 1919: 38: synonymy by Muir and Giffard, 1924: 31.</li> <li>= <i>Liburnia albicollis</i> Haupt, 1935 [nec. Motschulsky 1863]: 144; synonymy by Metcalf, 1943: 498; see also Wagner 1954: 217-8 (synonymy with <i>Liburnia albicollis</i> Motschulsky 1863, e.g., Lindberg, 1936: 18; China 1938: 54, Linnavouri 1952: 191, is in error).</li> <li>= <i>Delphacodes shirozui</i> Ishihara, 1949: 53-54: synonymy by Fennah, 1956: 122.</li> <li>= <i>Calligypona propinqua</i> (Fieber): new combination by Dlabola, 1954: 14, 64 (combination by implication in China 1954: 165).</li> <li>= <i>Toya propinqua</i> (Fieber): new combination by Fennah, 1964: 142 (from <i>Delphacodes propinqua</i>); see also Fennah 1965b: 56.</li> <li>= <i>Metadelphax propinqua</i> (Fieber): new combination by Wagner 1963: 170; synonymy by implication Fennah 1964: 142 (as <i>Toya propinqua</i>); restored combination by Ding 2006: 511, 513-514; Gonzon and Bartlett, 2008: 242-246.</li> <li>= <i>Calligypona (Metadelphax) propinqua</i> (Fieber): new combination by Linnavuori, 1964: 341-342.</li> <li>= <i>M[etadelphax] propinqua</i> (Fieber): misspelling by Mitjaev, 1971: 66.</li> <li>= <i>Toya (Calligypona) propinqua</i> (Fieber): new combination by Linnavouri, 1973: 107.</li> <li>= <i>Toya propinqua neopropinqua</i> (Muir): new status by Kuoh et al 1983: 155.</li> <li>= <i>Toya porpinqua</i> (Fieber): misspelling by Yang, 1989: 219.</li> </ul>	<p>Panama, Paraguay, Puerto Rico (inc. Vieques Is., Caja de Muertos Is.), Peru, St. Lucia, Suriname, Trinidad and Tobago, Venezuela; Virgin Islands (St. Croix, St. Thomas, St. Vincent).</p> <p>OLD WORLD: Australia, Cyprus, Egypt, Ethiopia, India, Israel, Ivory Coast, Japan, Kenya, Kyrgyzstan, Libya, Madagascar, Morocco, Niger, Northern Mariana Islands, Palau, Rhodesia, Thailand, Turkey, United Arab Emirates, Uzbekistan, Zambia, Zimbabwe. Widespread in Palearctic (e.g. Nast 1972, 1987; Holzinger et al. 2003); Afrotropics (e.g. Fennah 1969a, Asche 1988), Australia (Fennah 1965b), and Indomalayan (e.g., Ishihara 1949, Lee 1979, Yang 1989, Ding 2006) regions.</p>
---	---

<p>= <i>Toya propinqua</i> (Fieber): misspelling by Rodriguez-León et al. 1994a: 19.</p>	
<p><i>wetmorei</i> (Muir &amp; Giffard, 1924)          = <i>Megamelus terminalis</i> Crawford, 1914: 623 [nec. Van Duzee 1907] [Unavailable, preoccupied].          = <i>Delphacodes wetmorei</i> Muir and Giffard, 1924: 30, fig. 33, 104 (Replacement name for unavailable <i>Megamelus terminalis</i> Crawford, 1914).          = <i>Metadelphax wetmorei</i> (Muir &amp; Giffard, 1924); new comb. by Gonzon and Bartlett, 2008: 246-248.</p>	<p>USA: FL, GA, LA, NC: ?Mexico (Nicaragua reported in error, see Gonzon and Bartlett 2008: 248)</p>
<p><b>Muellerianella</b> Wagner, 1963</p>	
<p><i>laminalis</i> (Van Duzee, 1897)          = <i>Liburnia laminalis</i> Van Duzee, 1897.          = <i>Delphacodes laminalis</i> (Van Duzee, 1897); comb. nov. by          = <i>Megamelus laminalis</i> (Van Duzee, 1897); comb. nov. by Crawford, 1914: 607, 624.          = <i>Muellerianella laminalis</i> (Van Duzee, 1897); comb. nov. by Maw et. al., 2000: 85.          = <i>Liburnia lateralis</i> Van Duzee, 1897 (Nec. <i>Liburnia striatella</i> var. <i>lateralis</i> Fieber 1879).          = <i>Delphacodes lateralis</i> (Van Duzee, 1897); Comb. by Muir and Giffard, 1924: 25; Syn. by Oman, 1947: 212-213.          = <i>Delphacodes laterana</i> Metcalf, 1943; nom. nov. for <i>Delphacodes lateralis</i> (Van Duzee, 1897) by Metcalf, 1943: 458; syn. by Oman, 1947: 212-213.</p>	<p>USA: CA, CT, DC, DE, FL, IL, IN, KY, LA, MA, MD, ME, MO, MS, NC, NH, NY, OH, PA, TN;          CAN: ON QC; Cuba (Rodriguez-Leon et al. 1994)</p>
<p><b>Neomegamelanus</b> McDermott, 1952</p>	
<p><i>elongatus reductus</i> (Caldwell, 1951)          = <i>Megamelanus elongatus reductus</i> Caldwell, 1951.          = <i>Neomegamelanus elongatus reductus</i> (Caldwell, 1951); new comb. by McDermott, 1952:50-52 (subspecies by implication).</p>	<p>Puerto Rico (inc. Vieques Island).</p>
<p><i>graminicola</i> (Muir, 1928)          = <i>Megamelanus graminicola</i> Muir, 1928.          = <i>Neomegamelanus graminicola</i> (Muir, 1928); new comb. by McDermott, 1952:52.</p>	<p>Bermuda</p>
<p><i>lautus</i> (Metcalf, 1923)          = <i>Megamelanus lautus</i> Metcalf, 1923.          = <i>Neomegamelanus lautus</i> (Metcalf, 1923) new comb. by McDermott, 1952:52-</p>	<p>USA: TX; Mexico.</p>

54.	
<b><i>Neoperkinsiella</i></b> Muir, 1926	
<i>guaduae</i> Muir, 1926	Ecuador
<i>testacea</i> (Fowler, 1905) = <i>Goniolcium testaceum</i> Fowler, 1905. = <i>Stobaera testaceum</i> (Fowler, 1905); new comb. by Muir 1913: 243 (as <i>S. testacia</i> ); also Crawford, 1914: 576. = <i>Neoperkinsiella williamsi</i> Muir, 1926; syn. by Kramer, 1973: 400-401. = <i>Neoperkinsiella testacea</i> (Fowler, 1905); new comb. by Kramer, 1973: 400-401.	Canal Zone, Ecuador, Guatemala
<b><i>Nicetor</i></b> Fennah, 1963	
<i>brunnea</i> (Muir, 1926) = <i>Dicranotropis brunnea</i> Muir, 1926. = <i>Nicetor brunnea</i> Muir, 1926; new comb. by Fennah, 1963: 700.	Argentina
<i>pallidinervis</i> (Muir, 1926) = <i>Dicranotropis pallidinervis</i> Muir, 1926 = <i>Nicetor pallidinervis</i> (Muir, 1926); new comb. by Fennah, 1963: 700.	Brazil
<b><i>Nilaparvata</i></b> Distant, 1906	
<i>caldwelli</i> Metcalf, 1955 = <i>Nilaparvata muii</i> Caldwell, 1951 (in Caldwell and Martorell 1951): 192-194 (nec. China, 1925). = <i>Nilaparvata caldwelli</i> Metcalf, 1955: 262 (replacement name for unavailable <i>N. muii</i> Caldwell). = <i>Nilaparvata bis</i> Nast 1984: 396 (replacement name for unavailable <i>N. muii</i> Caldwell).	USA: FL, NC; Belize, Hispaniola, Puerto Rico; also reported Senegal (Fennah 1961) and Sudan (Linnavuori 1973).
<i>gerhardi</i> (Metcalf, 1923) = <i>Liburnia gerhardi</i> Metcalf, 1923: 210, pls. 62 & 70. = <i>Euidella gerhardi</i> (Metcalf): Metcalf, 1949: 58, pl. 1 & 4. = <i>Euides gerhardi</i> (Metcalf): Metcalf, 1952: 230-231. = <i>Nilaparvata gerhardi</i> (Metcalf, 1923); new comb. by Bartlett: 2007: 55-57.	USA: IL, LA, NH, NJ, NC, VA; ?Guatemala.
<i>serrata</i> Caldwell, 1951	USA: FL; Belize, ?Bolivia, Cayman Islands, Costa Rica, Mexico, Nicaragua, Puerto Rico,
<i>wolcottii</i> Muir and Giffard, 1924	USA: AZ, CA, CO, DE, FL, MD, NJ, UT; Mexico; Puerto Rico.
<b><i>Nothodelphax</i></b> Fennah, 1963	
<i>atlanticus</i> (China, 1958) = <i>Delphacodes atlanticus</i> China, 1958. = <i>Nothodelphax atlanticus</i> (China, 1958): new	Chile

<p>comb. by Fennah, 1965: 253.  = <i>Nothodelphax atlanticus atlanticus</i> China, 1958; new status by Fennah, 1965: 253.</p>	
<p><i>atlanticus nigrescens</i> Fennah, 1965</p>	Chile (Isla Wellington)
<p><i>foveata subfoveata</i> (Muir &amp; Giffard, 1924)  = <i>Delphacodes foveala subfoveata</i> Muir and Giffard, 1924.  = <i>Nothodelphax foveata subfoveata</i> (Muir and Giffard, 1924); new comb. by Fennah, 1963: 15.</p>	Mexico
<p><i>occlusa</i> (Van Duzee, 1897)  = <i>Liburnia occlusa</i> Van Duzee, 1897.  = <i>Megamelus occlusus</i> (Van Duzee, 1897); new comb. by Crawford, 1914: 619.  = <i>Delphacodes occlusa</i> (Van Duzee, 1897); new comb. by Muir and Giffard, 1924: 21.  = <i>Nothodelphax occlusa</i> (Van Duzee, 1897); new comb. by Fennah, 1965: 254.</p>	Mexico
<p><i>slossonae</i> (Ball, 1903)  = <i>Liburnia slossoni</i> Ball, 1903.  = <i>Megamehis slossoni</i> (Ball, 1903); new comb. by Crawford, 1914 :631.  = <i>Chloriona slossoni</i> (Ball, 1903); new comb. by Muir and Giffard, 1924: 8.  = <i>Chloriona slossonae</i> (Ball, 1903); emendation by Caldwell and Martorell, 1951: 160-161 (not listed as new)  = <i>Nothodelphax slossonae</i> (Ball, 1903); new comb. by Wilson and Hilburn, 1991: 416 (as <i>N. slossoni</i>).</p>	USA; FL; Bermuda, Cuba, Puerto Rico,
<p><b><i>Opiconsiva</i></b> Distant, 1917</p>	
<p><i>dodona</i> (Fennah, 1965)  = <i>Corbula dodona</i> Fennah, 1965  = <i>Opiconsiva dodona</i> (Fennah, 1965); new comb. by Fennah, 1975: 112.</p>	?Jamaica (Pacific & SE Asia)
<p><b><i>Peregrinus</i></b> Kirkaldy, 1904  = <i>Perigrinus</i> Kirkaldy, 1904; Missp. By Barber, 1914: 528, Van Der Goot, 1936: 25; Beamer, 1948: 106.</p>	
<p><i>maidis</i> (Ashmead, 1890)  = <i>Delphax maidis</i> Ashmead, 1890.  = <i>Dicranotropis maidis</i> (Ashmead, 1890); new comb. by Van Duzee, 1897: 227, 240.  = <i>Delphax psylloides</i> Lethierry, 1894; syn. by Kirkaldy, 1907: 132.  = <i>Liburnia psylloides</i> (Lethierry, 1894); comb</p>	USA: AL, DC, FL, HI, IL, LA, NC, OH, SC, TN, TX; Antigua, Argentina, Barbados, Bermuda, Brazil, Cayman Is. (Grand Cayman), Costa Rica, Cuba, Dominica, Haiti, Grenada, Jamaica, Mexico, Nevis, Nicaragua, Puerto Rico (inc. Vieques Island), St. Croix, St. Kitts, St. Lucia, St. Vincent, Trinidad

<p>by Melichar, 1903: 101.  = <i>Pundaluoya simplicia</i> Distant, 1906; Syn by Kirkaldy, 1907: 132.  = <i>Megamelus teapae albonotatus</i> Crawford, 1914; Syn. by Beamer, 1948: 106.  = <i>Delphacodes albonotata</i> (Crawford, 1914) (Nec. Muir and Giffard, 1924); comb. by Muir and Giffard, 1924: 36.  = <i>Perigrinus</i> (sic) <i>mardis</i> (Ashmead, 1890) missp. By Beamer, 1948: 106.</p>	<p>Canary Is., Dutch East Indies, Fiji, Galapagos, Guam, India, Indonesia (Java, Ambon, Borneo), Malay Peninsula, Mentawai Islands, New Caledonia and Loyalty Islands; Philippine Islands (Luzon), Sri Lanka, Tahiti, Taiwan,   Australia (Queensland, New South Wales)   Cape Province, Gold Coast, Mauritius (Rodrigues Island), Natal, Nigeria, Rhodesia, Seychelles, Sierra Leone, South Africa, Tanganyika</p>
<b><i>Perkinsiella</i> Kirkaldy, 1903</b>	
<b><i>saccharicida</i> Kirkaldy, 1903</b>	USA: FL, HI, LA; Fiji, Mauritius
<b><i>Phrictopyga</i> Caldwell, 1951 (in Caldwell and Martorell, 1951)</b>	
<p><i>contorta</i> (Muir, 1926)  = <i>Kelisia contorta</i> Muir, 1926.  = <i>Phrictopyga contorta</i> (Muir, 1926); new comb. by Caldwell and Martorell, 1951: 171.  = <i>Sogata parvula</i> Osborn, 1926; Syn. By Caldwell and Martorell, 1951: 171.</p>	Brazil, Jamaica, Puerto Rico
<p><i>curvistilus</i> (Muir, 1926)  = <i>Kelisia curvistilus</i> Muir, 1926.  = <i>Phrictopyga curvistilus</i> (Muir, 1926); new comb. by Caldwell and Martorell, 1951: 171.</p>	Brazil
<p><i>escadensis</i> (Muir, 1926)  = <i>Kelisia escadensis</i> Muir, 1926.  = <i>Phrictopyga escadensis</i> (Muir, 1926); new comb. by Caldwell and Martorell, 1951: 171.</p>	Brazil
<p><i>fuscovittata</i> (Muir, 1926)  = <i>Kelisia fuscovittata</i> Muir, 1926.  = <i>Phrictopyga fuscovittata</i> (Muir, 1926); new comb. by Caldwell and Martorell, 1951: 171.</p>	Brazil
<i>graminicola</i> (Muir, 1926)	Brazil
<i>holmgreni</i> (Muir, 1930)	Bolivia
<i>nugax</i> Fennah, 1959	Trinidad (St. Augustine)
<p><i>occidentalis</i> (Muir, 1926)  = <i>Kelisia occidentalis</i> Muir, 1926.  = <i>Phrictopyga occidentalis</i> (Muir, 1926); new comb. by Caldwell and Martorell, 1951: 171-172.  = <i>Sogata aurantii</i> Osborn, 1938 (Nec. Crawford, 1914); syn. by Caldwell and Martorell, 1951: 171-172.</p>	USA: TX, Ecuador, Mexico, Puerto Rico

<i>semele</i> Fennah, 1959	Trinidad (St. Augustine)
<i>urbana</i> (Muir, 1926) = <i>Kelisia urbana</i> Muir, 1926. = <i>Phrictopyga urbana</i> (Muir, 1926); new comb. by Caldwell and Martorell, 1951: 171.	Brazil
<b><i>Pissonotus</i></b> Van Duzee, 1897 = <i>Phyllodictus</i> Ball, 1926; syn. by Morgan and Beamer, 1949: 97; Bartlett and Deitz, 2000: 13.	
<i>abdominalis</i> (Crawford, 1914) = <i>Dicranotropis bakeri abdominalis</i> Crawford, 1914. = <i>Pissonotus abdominalis</i> (Crawford, 1914); new comb. by Bartlett and Deitz, 2000: 102.	Belize, Costa Rica, Dominican Republic, Honduras, Mexico, Nicaragua.
<i>absenta</i> Caldwell, 1951 <i>incertae sedis</i> by Bartlett & Deitz, 2000: 146	Puerto Rico
<i>albivultus</i> Morgan and Beamer, 1949	USA: FL, LA, ?TX; Belize, Costa Rica, Guatemala, Honduras, Mexico, Nicaragua, ?Venezuela.
<i>albovenosus</i> Osborn, 1935 = <i>Prokelisia albovenosus</i> Osborn, 1935; lapsus by McCoy and Rey, 1981; 289.	USA: AL, AZ, AR, CA, CT, DE, FL, GA, IL, LA, MD, MA, ?MO, MS, NH, NJ, NM, NY, NC, SC, TX, VA; Dominican Republic, Jamaica, Mexico, Puerto Rico.
<i>binotatus</i> Spooner, 1912 = <i>Dicranotropis binotatus</i> (Spooner, 1912); new comb by Crawford, 1914: 601.	USA: AL, AR, DE, FL, GA, LA, MS, MO, NC, TN, TX, VA; Bermuda
<i>boliviensis</i> Bartlett, 2000	Argentina, Bolivia, Brazil, Uruguay
<i>brevistilus</i> Bartlett, 2000	Mexico
<i>brunneus</i> Van Duzee, 1897 = <i>Pissonotus divaricatus</i> Spooner, 1912; syn. by Morgan and Beamer, 1949: 115-117. = <i>Dicranotropis divaricatus</i> (Spooner, 1912); new comb by Crawford, 1914: 600 = <i>Dicranotropis brunneus</i> (Spooner, 1912); new comb by Crawford, 1914: 594, 599.	USA: AL, AR, CT, DE, DC, FL, GA, IL, IN, IO, KS, KY, LA, MD, MA, MI, MS, MO, NH, NJ, NY, NC, OH, OK, PA, SC, TN, TX, VA, WI, WV; Canada: Nova Scotia, Ontario, Quebec, Alberta; Mexico (?Jamaica record in error; specimen label referred to Jamaica on Long Island in New York)
<i>decussatus</i> Bartlett, 2000	Guatemala, Mexico
<i>delicatus</i> Van Duzee, 1897 = <i>Pissonotus delicatus delicatus</i> Van Duzee, 1897; subspecific status implied by description of <i>P. d. melanurus</i> Van Duzee, 1917. = <i>Pissonotus pallipes</i> Van Duzee, 1897; syn. by Oman, 1947: 217-218, 220. = <i>Dicranotropis delicatus</i> (Van Duzee, 1897); new comb by Crawford, 1914: 594, 596-597.	USA: AL, AZ, AR, CA, CO, FL, GA, ID, IL, IO, KS, MS, MO, MT, NJ, NM, NY, NC, ND, OH, OK, OR, TX, UT, VT, WA, WY; Canada: Alberta, British Columbia, Saskatchewan, Quebec; Mexico, ?Jamaica

<p>= <i>Dicranotropis pallipes</i> (Van Duzee, 1897); new comb by Crawford, 1914: 594, 599.</p> <p>= <i>Pissonotus giffardi</i> Van Duzee 1925: syn. by Morgan and Beamer, 1949.</p>	
<i>dentatus</i> Morgan and Beamer, 1949	USA: FL, IL, NC; Cuba
<i>festucae</i> Bartlett, 2000	USA: AZ?; Mexico, ?Puerto Rico
<i>haywardi</i> Muir, 1929	Argentina, Brazil
<i>incertae sedis</i> by Bartlett & Deitz, 2000: 146.	
<i>jamaicensis</i> Bartlett, 2000	Jamaica
<p><i>muii</i> Metcalf, 1943</p> <p>= <i>Megamelus albicollis</i> Crawford, 1914 (nec. Motschulsky, 1863).</p> <p>= <i>Pissonotus albicollis</i> (Crawford, 1914): comb. by Muir and Giffard, 1924, 11.</p> <p>= <i>Pissonotus muii</i> Metcalf, 1943; Nom. Nov. for unavailable <i>M. albicollis</i> Crawford, 1914.</p>	Belize, Costa Rica, El Salvador, Guatemala, Honduras, Nicaragua, Panama
<p><i>neotropicus</i> (Muir, 1926)</p> <p>= <i>Phyllodinus neotropicus</i> Muir, 1926: 18-19.</p> <p>= <i>Phyllodictus neotropicus</i> (Muir, 1926); comb. by Metcalf, 1943: 146.</p> <p>= <i>Pissonotus neotropicus</i> (Muir, 1926); comb. by Bartlett &amp; Deitz, 2000: 64-66.</p>	Argentina, Bolivia, Brazil, Ecuador, Guyana, Paraguay, Peru, Venezuela
<p><i>nitens</i> (Van Duzee, 1909)</p> <p>= <i>Phyllodinus nitens</i> Van Duzee, 1909.</p> <p>= <i>Phyllodictus nitens</i> (Van Duzee, 1909); comb. by Ball, 1926.</p> <p>= <i>Pissonotus nitens</i> (Van Duzee, 1909); comb. by Morgan and Beamer 1949: 100, 107-108.</p>	USA: AL, FL, MS; Jamaica.
<i>paraguayensis</i> Bartlett, 2000	Brazil, ?Bolivia, Paraguay
<p><i>piceus</i> (Van Duzee, 1894)</p> <p>= <i>Megamelus piceus</i> Van Duzee, 1894.</p> <p>= <i>Dicranotropis bakeri</i> Crawford, 1914; syn. by Bartlett &amp; Deitz, 2000: 107-116.</p> <p>= <i>Pissonotus fulvus</i> Metcalf, 1923; syn. by Morgan and Beamer, 1949: 137-138.</p> <p>= <i>Delphacodes pictifrons</i> Osborn, 1938; syn. by Bartlett &amp; Deitz, 2000: 107-116.</p> <p>= <i>Pissonotus piceus</i> (Van Duzee, 1894); comb. by Oman, 1947: 219-220.</p>	USA: AL, AR, AZ, CT, DE, DC, FL, GA, IL, IN, KS, LA, MD, MA, MI, MO, MS, MT, NC, NH, NJ, NY, OH, PA, SC, TN, TX, VT, VA, WV; Canada; Ontario; Belize, Bermuda, Costa Rica, Cuba, Honduras, Jamaica, Mexico, Nicaragua, Panama, Peru, Trinidad, Venezuela.
<p><i>quadripustulatus</i> (Van Duzee, 1909)</p> <p>= <i>Stobaera 4-pustulata</i> Van Duzee, 1909.</p> <p>= <i>Pissonotus foveatus</i> Spooner 1912; syn. by Van Duzee, 1916: 84.</p> <p>= <i>Pissonotus variegatus</i> Spooner, 1912; syn. by Metcalf, 1923: 148.</p> <p>= <i>Dicranotropis varitgatus</i> (Spooner 1912); comb. by Crawford, 1914: 600.</p>	USA: AL, FL, GA, LA, TX; Mexico

<p>= <i>Dicranotropis foveatus</i> (Spooner, 1912); comb. by Crawford, 1914: 600.</p> <p>= <i>Stobaera quadripustulata</i> Van Duzee, 1909; transliteration of <i>4-pustulata</i> by Crawford, 1914: 572, 576.</p> <p>= <i>Stobaera foveatus</i> (Spooner, 1912); comb. by Van Duzee, 1916: 84.</p> <p>= <i>Pissonotus quadripustulatus</i> (Van Duzee, 1909); comb. by Metcalf, 1923: 148, 170.</p> <p>= <i>Pissonotus 4-pustulata</i> (Van Duzee, 1909); comb. by Morgan and Beamer, 1949: 110, 120-121.</p> <p>= <i>Pissonotus quadripustulatus</i> (Van Duzee, 1909); missp. By Stiling, 1994.</p>	
<i>spatulatus</i> Bartlett, 2000	USA: CA; Mexico
<b><i>Pygospina</i></b> Caldwell, 1951 (in Caldwell and Martorell, 1951)	
<p><i>aurantii</i> (Crawford, 1914)</p> <p>= <i>Megamelus aurantii</i> Crawford, 1914.</p> <p>= <i>Stenocranus hinei</i> Dozier, 1922; syn. by Muir, 1926: 25.</p> <p>= <i>Sogata aurantii</i> (Crawford, 1914); comb. by Muir and Giffard, 1924: 16.</p> <p>= <i>Pygospina aurantii</i> (Crawford, 1914); new comb. by Caldwell and Martorell, 1915: 169.</p>	Brazil, Cuba, Ecuador, Guatemala, Guyana, Puerto Rico
<i>reducta</i> Caldwell, 1951	Puerto Rico
<p><i>rezendensis</i> (Muir, 1926)</p> <p>= <i>Kelisia rezendensis</i> Muir, 1926</p> <p>= <i>Pygospina rezendensis</i> (Muir, 1926); new comb. by Caldwell and Martorell, 1951: 169.</p>	Brazil
<i>spinata</i> Caldwell, 1951	USA: FL; Cuba, Puerto Rico
<p><i>spinigera</i> (Fennah, 1945)</p> <p>= <i>Delphacodes spinifera</i> Fennah, 1945</p> <p>= <i>Pygospina spinigera</i> (Fennah, 1945); new comb. by Caldwell and Martorell, 1951: 169.</p>	Trinidad
<b><i>Pyrophagus</i></b> Remes Lenicov & Varela, 2005 (cited by Velazquez et al. 2006). (Note: this taxon and its species may be nomen nud. – description cited as ‘inédito’, have not found published description)	
<p><i>tigrinus</i> Remes Lenicov &amp; Varela, 2005 (cited by Velazquez et al. 2006).</p> <p>= <i>Delphacodes tigrinus</i> Varela and Varela, 1999 (<i>nomen nud.</i>)</p>	Argentina
<b><i>Sogatella</i></b> Fennah, 1956	
= <i>Sogatodes</i> Fennah, 1963; syn. by Asche and	

<p>Wilson, 1990; 5. = <i>Chloriona (Sogatella)</i> Fennah, 1956.</p>	
<p><i>furcifera</i> (Horváth, 1899) = <i>Delphax furcifera</i> Horváth, 1899. = <i>Liburnia furcifera</i> (Horváth, 1899); comb. by Matsumura, 1900: 262. = <i>Megamelus furcifera</i> (Horváth, 1899); comb. by Muir, 1917. = <i>Liburnia (Delphax) furcifera</i> (Horváth, 1899); comb. by Kuwana, 1930. = <i>Liburnia albolineosa</i> Fowler, 1905: syn. by Caresche, 1933: 492. = <i>Megamelus albolineosa</i> (Fowler, 1905); comb. by Crawford, 1914: 610. = <i>Sogata distincta</i> Distant, 1912: syn. by Muir, 1919: 7. = <i>Sogata pallescens</i> Distant, 1916: syn. by Caresche, 1933: 492, 498. = <i>Opiconsiva colorata</i> Distant, 1917: syn. by Muir and Giffard, 1924: 15. = <i>Opiconsiva derelicta</i> Distant, 1917: syn. by Muir and Giffard, 1924: 15; removed from syn. by Asche and Wilson, 1990: 7 (to <i>Sogatella kolophon</i>). = <i>Opiconsiva balteata</i> Distant, 1917: syn. by Muir and Giffard, 1924: 15; removed from syn. by Asche and Wilson, 1990: 7 (to <i>Sogatella kolophon</i>). = <i>Opiconsiva gloriosus</i> Distant, 1917: syn. by Muir and Giffard, 1924: 15 (error); syn. with <i>Matutinus opulentus</i> Distant, 1917 by Fennah, 1963: 48. = <i>Delphax nigrigenis</i> Jacobi, 1917; syn. by Muir and Giffard, 1924: 15. = <i>Sogata kyusyuensis</i> Matsumura and Ishihara, 1945; syn. by Asche and Wilson, 1990: 5, 12.</p>	<p>Widespread in Paletropics; reported from Neotropics (e.g., Puerto Rico; Caldwell and Martorell, 1951, Trinidad; Fennah, 1945), but Asche and Wilson: 1990 could not verify the presence of this species in the New World, noting that “records of <i>S. furcifera</i> from the New World Counties (Muir and Giffard, 1924: 13; Muir 1926: 26) in most cases concern <i>S. molina</i> (Fennah) or <i>S. kolophon</i> (Kirkaldy).” The presence of <i>Sogatella furcifera</i> remains unverified in the New World.</p>
<p><i>kolophon</i> (Kirkaldy, 1907) = <i>Delphax kolophon</i> Kirkaldy, 1907. = <i>Sogatella kolophon meridiana</i> (Beamer, 1952); comb. by Fennah, 1963: 59. = <i>Delphacodes meridiana</i> Beamer, 1952; syn. by Asche and Wilson, 1990: 7, 16. = <i>Sogatella kolophon atlantica</i> Fennah, 1963; syn. by Asche and Wilson, 1990: 7, 16. = <i>Chloriona (Sogatella) kolophon</i> (Kirkaldy, 1907); new comb. by (? see Fennah, 1959). = <i>Opiconsiva insularis</i> Distant, 1917; syn. by Asche and Wilson, 1990: 7, 16.</p>	<p>Nearctic region: USA: CA, FL, GA, KY, MO, MS, TX; CAN: NS; Bermuda Islands  Palearctic region: Azores, Canary Islands, Japan, Korea  Afrotropical region: Cape Verde Island, Ivory Coast, Mauritius, Nigeria, Rodrigues Island, South Africa.  Indomalayan region: Cambodia, China, Hong Kong, India, Indonesia, Laos, Malaysia, Papua New Guinea, Philippine Islands, Seychelles Islands, Sri Lanka, Thailand, Vietnam</p>

<p>= <i>Sogatella kolophon insularis</i> Distant, 1917; comb. by Fennah, 1963: 59.</p> <p>= <i>Opiconsiva derelicta</i> Distant, 1917: syn. by Asche and Wilson, 1990: 7, 16.</p> <p>= <i>Sogatella derelicta</i> Distant, 1917; comb. by Fennah, 1963: 62.</p> <p>= <i>Opiconsiva balteata</i> Distant, 1917: syn. by Asche and Wilson, 1990: 7, 16.</p> <p>= <i>Sogatella balteata</i> (Distant, 1917); comb. by Fennah, 1963: 64.</p> <p>= <i>Sogatella chenhea</i> Kuoh, 1977: syn. by Asche and Wilson, 1990: 7, 16.</p> <p>= <i>Delphacodes elegantissima</i> Ishihara, 1952: syn. by Asche and Wilson, 1990: 7, 16.</p> <p>= <i>Sogatella elegantissima</i> (Ishihara, 1952); comb. by Fennah, 1963: 67.</p> <p>= <i>Sogatella nebris</i> Fennah, 1963: syn. by Asche and Wilson, 1990: 7, 17.</p>	<p>Pacific Region: Belau Island, Bonin Island, Eniwetok Atoll, Fiji, Galapagos, Guam, Hawaii, Mangareva Island, Marquesas Island, Marshall Island, Micronesia, New Caledonia, Northern Mariana Islands, Pitcairn Island, Society Islands, Solomon Islands, Tonga Islands, Western Samoa</p> <p>Australian region: Australia</p> <p>Neotropical region: Argentina, Ecuador, Guyana, Jamaica, Mexico, Montserrat, St. Lucia, Venezuela</p> <p>Atlantic Ocean: St. Helena Island.</p>
<p><i>molina</i> (Fennah, 1963)</p> <p>= <i>Sogatodes molinus</i> Fennah, 1963.</p> <p>= <i>Sogatella molina</i> (Fennah, 1963): new comb. by Asche and Wilson, 1990: 7.</p>	<p>USA: FL; Cayman Is. (Grand Cayman), Bermuda, Mexico</p>
<p><b><i>Sparnia</i></b> Stål, 1862</p>	
<p><i>edwardsi</i> Muir, 1927</p>	<p>Chile (Chiloe Is.)</p>
<p><i>praecellens</i> Stål, 1862</p>	<p>Brazil, Ecuador, Venezuela</p>
<p><i>williamsi</i> Muir, 1926</p>	<p>Ecuador</p>
<p><b><i>Stobaera</i></b> Stål, 1859</p> <p>= <i>Goniolcium</i> Fowler, 1905; syn. by Muir, 1915: 264.</p>	
<p><i>azteca</i> Muir, 1913</p>	<p>Guatemala, Mexico, Nicaragua</p>
<p><i>bahamensis</i> Metcalf, 1954</p>	<p>Bahamas</p>
<p><i>caldwelli</i> Kramer, 1973</p>	<p>USA: AZ, CA, NM, NV, TX, UT; Mexico</p>
<p><i>concinna</i> (Stål, 1854)</p> <p>= <i>Delphax concinna</i> Stål, 1854.</p> <p>= <i>Stobaera concinna</i> (Stål, 1854); new comb. by Stål: 1859: 327.</p> <p>= <i>Stobaera minuta</i> Osborn, 1905; syn. by Kramer, 1973: 388-391.</p>	<p>USA: AZ, CA, CO, FL, ?IL, LA, ?MS, ?NJ, TX, UT; Bermuda, Cuba, Hispaniola, Mexico, ?St. Lucia</p>
<p><i>granulosa</i> (Fowler, 1905)</p> <p>= <i>Goniolcium granulosa</i> Fowler, 1905.</p> <p>= <i>Stobaera granulosa</i> (Fowler, 1905); comb. by Muir, 1913: 242.</p> <p>= <i>Stobaera granulosum</i> (Fowler, 1905); unjustified emendation by Crawford, 1914: 576. .</p>	<p>Ecuador, Guatemala, Mexico.</p>
<p><i>koebeli</i> Muir, 1913</p>	<p>Guatemala, Mexico</p>
<p><i>pallida</i> Osborn, 1905</p>	<p>USA: DE, FL, MD, NY, VA; Mexico</p>
<p><i>tricarinata</i> (Say, 1825)</p> <p>= <i>Delphax tricarinata</i> Say, 1825.</p>	<p>USA: AL, AZ, CA, CO, DE, IA, ID, FL, GA, ID, IL, IN, IO, KS, LA, MA, MD, MN, MO, MS, NB,</p>

<p>= <i>Stobaera tricarinata</i> (Say, 1925); comb. by Osborn, 1938: 239.  = <i>Stobaera nigripennis</i> Crawford, 1914; Syn by Kramer, 1973: 393-396.</p>	<p>NC, ND, NJ, NM, NV, NY, OH, OR, PA, SC, TN, TX, UT, VA, ?Guatemala, Mexico, ?Nicaragua, Cuba, Puerto Rico. (Neotropical records need confirmation)</p>
<p><b>Syndelphax</b> Fennah, 1963</p>	
<p><i>dissipatus</i> (Muir, 1926)</p>	<p>Brazil, Ecuador</p>
<p><i>dolosa</i> (Muir, 1926)  = <i>Delphacodes dolosa</i> Muir, 1926; new comb. by Gonzon and Bartlett, 2008: 252.</p>	<p>Brazil</p>
<p><i>disonymus</i> (Kirkaldy, 1907)  = <i>Delphax disonymos</i> Kirkaldy, 1907.  = <i>Delphacodes disonymos</i> (Kirkaldy, 1907), new comb. by Muir, 1920: 139.  = <i>Syndelphax disonymos</i> (Kirkaldy, 1907); new comb. by Fennah, 1963: 15-16.  = <i>Delphax matanitu</i> Kirkaldy, 1907: syn. by Muir 1920: 139; Fennah, 1975: 109.</p>	<p>USA: HI (Asche, 2000); Cayman Is. (Grand Caymen) (Reported by Fennah, 1971: 321 as <i>S. matanitu</i>).  Australia, Ceylon, Fiji, New Caledonia, South Africa, Western Pacific.</p>
<p><i>fallax</i> (Muir, 1926)  = <i>Delphacodes fallax</i> Muir, 1926  = <i>Syndelphax fallax</i> (Muir, 1926); new comb. by Gonzon and Bartlett, 2008: 252.</p>	<p>USA:TX; Brazil</p>
<p><b>Synpteron</b> Muir, 1926</p>	
<p><i>brazilensis</i> Muir, 1926</p>	<p>Brazil</p>
<p><b>Tagosodes</b> Asche and Wilson, 1990</p>	
<p><i>albifacies</i> (Caldwell, 1951)  = <i>Sogata albifacies</i> Caldwell, 1951.  = <i>Tagosodes albifacies</i> (Caldwell, 1951); new comb. by Asche and Wilson, 1990: 32.</p>	<p>USA: FL; Puerto Rico</p>
<p><i>albolineosus</i> (Fowler, 1905)  = <i>Liburnia albolineosus</i> Fowler, 1905.  = <i>Sogatodes albolineosus</i> (Fowler, 1905); syn. by Fennah, 1963: 74-75.  = <i>Tagosodes albolineosus</i> (Fowler, 1905); new comb. by Asche and Wilson, 1990: 32.</p>	<p>Cuba, ?Jamaica, Mexico, Puerto Rico, Nicaragua</p>
<p><i>anomalus</i> (Muir, 1926)  = <i>Sogata anomala</i> Muir, 1926.  = <i>Sogatodes anomalus</i> (Muir, 1926); new comb. by Fennah, 1963: 74.  = <i>Tagosodes anomalus</i> (Muir, 1926); new comb. by Asche and Wilson, 1990: 32.</p>	<p>Brazil</p>
<p><i>approximatus</i> (Crawford, 1914)  = <i>Megamelus approximatus</i> Crawford, 1914.  = <i>Sogata approximata</i> (Crawford, 1914); new comb. by Wolcott, 1923: 273.  = <i>Sogatodes approximatus</i> (Crawford, 1914); new comb. by Fennah, 1963: 74.  = <i>Chloriona (Sogatella) approximata</i></p>	<p>Nicaragua, Puerto Rico, Peru</p>

<p>(Crawford, 1914); new comb. by Fennah, 1959: 259.  = <i>Tagosodes approximatus</i> (Crawford, 1914); new comb. by Asche and Wilson, 1990: 32.</p>	
<p><i>biangulatus</i> (Muir, 1929)  = <i>Sogata biangulata</i> Muir, 1929.  = <i>Sogatodes biangulata</i> (Muir, 1929); new comb. by Fennah, 1963: 74.  = <i>Tagosodes biangulatus</i> (Muir, 1929); new comb. by Asche and Wilson, 1990: 32.</p>	Argentina
<p><i>cubanus</i> (Crawford, 1914)  = <i>Dicranotropis cubanus</i> Crawford, 1914.  = <i>Dicranotropis cubana</i> Crawford, 1914; emendation by Wiess, 1917: 215.  = <i>Megamelus flavolineatus</i> Muir, 1920; syn. by Wolcott, 1923: 273.  = <i>Sogata cubanus</i> (Crawford, 1914); comb. by Wolcott, 1923: 273.  = <i>Sogata cubana</i> (Crawford, 1914); emendation by Muir, 1926: 26-28.  = <i>Peregrinus cubana</i> (Crawford, 1914); Osborn, 1926: 358.  = <i>Liburnia cubana</i> (Crawford, 1914); new comb. by Metcalf, 1943: 354.  = <i>Chloriona (Sogatella) cubana</i> (Crawford, 1914); new comb. by Fennah, 1959: 259.  = <i>Sogatodes cubanus</i> (Crawford, 1914); new comb. by Fennah, 1963: 74.  = <i>Delphacodes pallidivitta</i> Fennah, 1945; syn. by Asche and Wilson, 1990: 33.  = <i>Chloriona (Sogatella) panda</i> Fennah, 1958; syn. by Fennah, 1969: 56.</p>	<p>USA: FL,NJ; Cuba, Puerto Rico, St. Lucia, Bermuda, Trinidad, Venezuela</p> <p>Nigeria, Ivory Coast</p>
<p><i>nauticus</i> (Muir, 1926)  = <i>Sogata nautica</i> Muir, 1926.  = <i>Chloriona (Sogatella) nautica</i> (Muir, 1926); new comb. by Fennah, 1959: 259.  = <i>Sogatodes nauticus</i> (Muir, 1926); new comb. by Fennah, 1963: 74.  = <i>Tagosodes nauticus</i> (Muir, 1926); new comb. by Asche and Wilson, 1990: 33.</p>	Guyana, Venezuela
<p><i>orizicolus</i> (Muir, 1926)  = <i>Sogata orizicola</i> Muir, 1926.  = <i>Sogatodes orizicola</i> (Muir, 1926); comb. by Fennah, 1963: 71.  = <i>Chloriona (Sogatella) orizicola</i> (Muir, 1926); new comb. by Fennah, 1959:</p>	<p>USA: FL, LA, MS, Brazil, Cayman Islands (Grand Cayman), Cuba, Guyana, Mexico, Puerto Rico, Dominica, Trinidad, Venezuela</p>

<p>259.  = <i>Sogatodes brazilensis</i> (Muir, 1926); new comb. by Fennah, 1963: 74.  = <i>Sogata brazilensis</i> Muir, 1926; syn. by Fennah, 1965: 215-217.</p>	
<p><i>wallacei</i> (Muir &amp; Giffard, 1924)  = <i>Sogata wallacei</i> Muir and Giffard, 1924.  = <i>Chloriona (Sogatella) wallacei</i> (Muir &amp; Giffard, 1924); new comb. by Fennah, 1959: 259.  = <i>Sogatella wallacei</i> (Muir and Giffard, 1924); new status by Fennah, 1963.  = <i>Tagosodes wallacei</i> (Muir and Giffard, 1924); new comb. by Asche and Wilson, 1990: 7, 36.</p>	<p>Belize, Cuba, Dominica, Guyana, Honduras, Jamaica, Puerto Rico</p>
<p><b>Toya</b> Distant, 1906  = <i>Himeunka</i> Matsumura and Ishihara 1945: syn. by Anufriev 1977: 865. Removed from synonymy by Kuoh et al. 1981: 190.  = <i>Metadelphax</i> Wagner, 1963: syn. by Fennah 1964: 142 (by implication; type species moved to <i>Toya</i>); Nast 1972: 65, Linnavuori 1973: 107 (generic synonymy listed). Removed from synonymy by Ding 2006: 511.</p>	
<p><i>balboae</i> (Muir and Giffard, 1924)</p>	<p>Argentina</p>
<p><i>boxi</i> (Muir &amp; Giffard, 1924)  = <i>Delphacodes boxi</i> Muir, 1926.  = <i>Toya boxi</i> (Muir &amp; Giffard, 1924); new combination by Fennah, 1965: 96.</p>	<p>USA: FL: British Guiana, Paraguay, Puerto Rico, St. Lucia, Trinidad and Tobago</p>
<p><i>dietrichi</i> Gonzon and Bartlett, 2008</p>	<p>Mexico (Colima)</p>
<p><i>iaxartes</i> (Fennah, 1959)  = <i>Delphacodes iaxartes</i> Fennah, 1959  = <i>Toya iaxartes</i> (Fennah, 1959); new combination by Fennah, 1965: 96.  <i>Nomen dubium</i> by Gonzon and Bartlett, 2008; 215-216, ill. 214.</p>	<p>USA: FL, LA, NC; Bahamas, Belize, Brazil, Bolivia, Cayman Islands, Colombia, Guatemala, Honduras, Panama, Paraguay, St. Lucia, Suriname, Uruguay</p>
<p><i>idonea</i> (Beamer, 1947)  = <i>Delphacodes idonea</i> Beamer, 1947.  = <i>Toya idonea</i> (Beamer, 1947); new comb. by Gonzon and Bartlett, 2008: 222-224.</p>	<p>USA: FL, LA, NC; Bahamas, Belize, Brazil, Bolivia, Cayman Is., Columbia, Guatemala, Honduras, Panama, Paraguay, Suriname, Uruguay, Venezuela</p>
<p><i>nigra</i> (Crawford, 1914)  = <i>Megamelus erectus niger</i> Crawford, 1914: 608, 624.  = <i>Delphacodes nigra</i> (Crawford, 1914), new combination and emendation by Muir and Giffard, 1924: 32.  = <i>Toya nigra</i> (Crawford, 1914); new comb. by Gonzon and Bartlett, 2008: 224-227.</p>	<p>USA: FL; Argentina, Belize, Bolivia, Brazil, British Guiana, Cayman Islands, Colombia, Costa Rica, Cuba, Dominica, Guadeloupe, Jamaica, Mexico, Panama, Puerto Rico (inc. Vieques Is.), Trinidad and Tobago, Venezuela</p>

<p>= <i>Delphacodes axonopi</i> Fennah, 1945: 434-435; syn. by Gonzon and Bartlett, 2008: 224.</p>	
<p><i>recurva</i> Gonzon and Bartlett, 2008</p>	<p>Mexico (Veracruz State).</p>
<p><i>venilia</i> (Fennah, 1959)          = <i>Delphacodes venilia</i> Fennah, 1959          = <i>Toya venilia</i> (Fennah, 1969); new comb. by Fennah, 1965: 96.</p>	<p>British Virgin Islands (Guana, Necker, Anegada, Beef, Great Camanoë, Great Dog, Great Thatch, Jost van Dyke, Tortola, Virgin Gorda); US Virgin Islands (St. John); Montserrat, Puerto Rico (incl. Muertos Is., Vieques Is.), Turks And Caicos Islands</p>

**Notes:**

A synonymy of *Bergias* with *Pentagramma* is listed by Ashe 1985: 283, but I found no prior listing of that synonymy; similarly that of *Micromasoria*, *Consualia*., and *Livatis* with *Ugyops*.

Ashe's unpublished list gives *Delphax producta* as a *Delphacodes*, but I have found no transfer of this species into *Delphacodes*.