

Potato Disease Advisory
May 13, 2009
Cooperative Extension System
University of Delaware

Bob Mulrooney, Extension Plant Pathologist

Late blight Advisory

If you would like a FAX or email report please call 302-831-4865, or email bobmul@udel.edu

We are using the E-WEATHER SERVICE from SkyBit, Inc as we have in the past. The service determines specific requested weather parameters (temperature, relative humidity and rainfall) based on calculations of data from the nearest National Weather Service stations. This weather data is used in the WISDOM software program for predicting late blight and making spray recommendations and accumulating p-days for early blight forecasting. Our location this year is:

Location: Shadybrook Farms, Little Creek, DE in Kent County.
Greenrow: May 1

Date	DSV	Total DSV	Spray Recommendation
5/3 -5/6		41	
5/7	2	43	
5/8	1	44	
5/9- 5/13	0	44	10- day spray interval

Disease severity values accumulated very quickly during the past rainy period. The threat of late blight from seed infection is low to medium, but there was some in Maine last season. Be vigilant, given this recent weather pattern. Apply sprays of mancozeb (Dithane, Manzate, Pencozeb, Manex II), Gavel or Bravo (chlorothalonil) now and again before plants canopy down the row. See the 2009 Commercial Vegetable Recommendations- Delaware for more information on fungicides.

Fortunately late blight has not been a problem here in Delaware for many years and unless you have seed from an unknown source the risk of late blight is low, but the weather was perfect for it if it was present on seed.